

HIV integrase^{R263A/K264A} is defective for TRN-SR2/IN interaction and nuclear import of the PIC

Stéphanie De Houwer, Jonas Demeulemeester, Wannes Thys, Zeger Debyser and Frauke Christ

Laboratory for Molecular Virology and Gene Therapy, Department of Pharmaceutical and Pharmacological Sciences, KU Leuven, Leuven, Belgium

Background: Transportin-SR2 (TRN-SR2/TNPO3) is a human karyopherin encoded by the *TNPO3* gene. We identified TRN-SR2 as a binding partner of HIV-1 integrase (IN) and validated TRN-SR2 as an important cellular cofactor for the nuclear import of HIV-1. However, the question still remained whether the direct interaction between TRN-SR2 and IN mediates the nuclear import of HIV.

Methods: By peptide screening we analyzed the interaction between TRN-SR2 and IN in molecular detail. Protein-protein interactions were verified by co-IP and AlphaScreen. We engineered site specific mutants in the C-terminal domain of HIV-1 IN that display reduced interaction with TRN-SR2 and studied their role in HIV replication using Q-PCR and fluorescence microscopy.

Results: Using the AlphaScreen protein-protein interaction assay we have been able to pinpoint the interacting hot spots in IN to R262/R263/K264 and K266/R269 in the IN C-terminal domain. We also identified a secondary interaction surface involving residues F185/K186/R187 and K188 in the catalytic core domain. Next, we introduced mutations at these positions in the C-terminal domain in the virus to corroborate the biological relevance of the interaction. Several mutations in the C-terminal domain of HIV IN inhibited the IN/TRN-SR2 interaction and rendered the virus replication-deficient. Some mutants affected reverse transcription (RT) compromising analysis of HIV nuclear import. All mutants also affected integrase activity. Still, one mutant, IN^{R263A/K264A}, retained full RT activity but displayed a specific block at the level of nuclear import as measured by Q-PCR and fluorescence microscopy. Although this mutant was defective for integration, no increase in 2-LTR circles was detected. Moreover, HIV encoding IN^{R263A/K264A} showed reduced nuclear import when assayed with an eGFP-IN labeled HIV.

Conclusion: The IN^{R263A/K264A} mutation in the C-terminal domain of HIV-1 integrase reduces the interaction with TRN-SR2 and specifically blocks HIV replication at the stage of nuclear import, corroborating the importance of this direct protein-protein interaction in HIV nuclear import.